



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

materials. Both scientific and non-scientific readers will find much to interest them in it. The illustrations, of which the book contains 143, are noteworthy.

H. H.

Contributions to the Geology of the Falkland Islands. By J. G. ANDERSSON. With 9 plates, and maps. Wissenschaftliche Ergebnisse der schwedischen Südpolar-Expedition, 1901-03, Band III, Lieferung 2. London: Dulau & Co.

In their deeply indented coast lines with numerous drowned valleys, the Falklands show a recent submergence of about 100 meters. During the ice age they stood at about their present height above the sea, while in pre-Glacial times they were somewhat higher and carried considerable rivers. A striking feature on East Falkland is the so-called "stone-rivers," which are level sheets of huge, angular boulders streaming down the hill-sides and reaching far out on almost level surfaces. This phenomenon is the product of solifluction, i. e., of gradual creep down the slopes of masses of waste saturated with water. Glacial action has not been the direct agent; for the islands seem never to have possessed a large ice cap. Thick peat deposits in this region furnish yet another instance of notable accumulations of vegetation in *cool*, moist climates. Devonian sandstone is found in the islands, resting on an Archean basement, while younger Paleozoic rocks are also present.

H. H.

The Meteor Crater of Canyon Diablo, Arizona; Its History, Origin, and Associated Meteoric Irons. By GEORGE P. MERRILL. Reprinted from Smithsonian Miscellaneous Collections (Quarterly Issue), Vol. L, Part 4, pp. 461-98, pls. LXI-LXXV, figs. 124-29. Washington, January 27, 1908.

The author inclines strongly to the view that the peculiar topographic feature commonly known as Coon Butte owes its origin to the impact of a meteorite of unprecedented size. The crater, which is 4,000 feet in diameter and 500 feet deep, lies in a region of undisturbed sedimentary rocks which are horizontally bedded except in the immediate vicinity of the crater itself, where they show a strong quaquaversal dip. Extensive development work, now being carried on by a mining company in the field, fails to substantiate the theory that volcanic action has been the factor involved, but shows that the disturbance was essentially superficial. Microscopic and megascopic studies of the fragmental materials in and about the crater indi-